

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification 5 :</b> A01N 57/20, 37/00 // (A01N 37/00 A01N 37:02, 37:06)	<b>A1</b>	<b>(11) International Publication Number:</b> WO 92/11764 <b>(43) International Publication Date:</b> 23 July 1992 (23.07.92)
<b>(21) International Application Number:</b> PCT/US91/09621 <b>(22) International Filing Date:</b> 20 December 1991 (20.12.91) <b>(30) Priority data:</b> 638,590 8 January 1991 (08.01.91) US <b>(71) Applicant:</b> MONSANTO COMPANY [US/US]; 800 North Lindbergh Boulevard, St. Louis, MO 63167 (US). <b>(72) Inventors:</b> ARNOLD, Kristin, Anne ; 1012 North Harrison Avenue, Kirkwood, MO 63122 (US). BUGG, Marvin, Wayne ; 1521 Virginia Avenue, Ellisville, MO 63011 (US). WHITE, Randall, Joseph ; 1823 Robin Hood Drive, Miamisburg, OH 45342 (US).		<b>(74) Agent:</b> BOLDING, James, Clifton; Monsanto Company, 800 North Lindbergh Boulevard, St. Louis, MO 63167 (US). <b>(81) Designated States:</b> AT (European patent), AU, BB, BE (European patent), BF (OAPI patent), BG, BJ (OAPI patent), BR, CA, CF (OAPI patent), CG (OAPI patent), CH (European patent), CI (OAPI patent), CM (OAPI patent), CS, DE (European patent), DK (European pa- tent), ES (European patent), FI, FR (European patent), GA (OAPI patent), GB (European patent), GN (OAPI patent), GR (European patent), HU, IT (European pa- tent), JP, KP, KR, LK, LU (European patent), MC (Eu- ropean patent), MG, ML (OAPI patent), MN, MR (OA- PI patent), MW, NL (European patent), NO, PL, RO, SD, SE (European patent), SN (OAPI patent), SU ,TD (OAPI patent), TG (OAPI patent).  <b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> IMPROVED HERBICIDAL FORMULATION  <b>(57) Abstract</b>  This invention relates to a novel aqueous agriculturally acceptable formulation, a process used to prepare it and a pesticid- al method of using it in application to plants or weeds to kill or control plants or weeds. The formulation comprises an effective amount of N-phosphonomethylglycine or an agriculturally acceptable salt of N-phosphonomethylglycine and an effective amount of at least one C <sub>5</sub> to C <sub>16</sub> or preferably a C <sub>8</sub> to C <sub>12</sub> agriculturally acceptable fatty acid itself or in the form of an agri- culturally acceptable water soluble salt or mixtures thereof. Other ingredients are optional, including surfactant(s), anti- foam(s) and antimicrobial(s) or other ingredients such as pesticides including herbicides, insecticides and fungicides.		

*Abn after BPAI Affid Exr.*

*Patents from this family*

*✓ 5,994,269 - related*

*09/264,589 → ⊕*

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MC	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	RU	Russian Federation
CG	Congo	KP	Democratic People's Republic of Korea	SD	Sudan
CH	Switzerland	KR	Republic of Korea	SE	Sweden
CI	Côte d'Ivoire	LI	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
DE	Germany	MC	Monaco	TG	Togo
DK	Denmark			US	United States of America

-1-

IMPROVED HERBICIDAL FORMULATIONField of the Invention

This invention relates to a novel aqueous  
5 agriculturally acceptable herbicidal formulation, a  
process used to prepare it and a herbicidal method of  
using it in applying it to weeds or plants to kill and  
control weeds or plants.

10 The formulation, preferably a herbicidally  
efficacious formulation, comprises an effective amount  
of an agriculturally acceptable salt of N-phosphono-  
methylglycine and a herbicidally effective amount of at  
least one agriculturally acceptable fatty acid or a salt  
15 thereof or a mixture of fatty acids and a salt(s)  
thereof or a mixture of salts of a fatty acid(s) giving  
in an aqueous solution to be applied to plants or a  
desired locus a pH in a desired range. Other  
ingredients are optional, including surfactant(s),  
20 antifoam(s), antimicrobial(s) or one or more additional  
pesticides including herbicides, insecticides and  
fungicides. As used herein, the term "agriculturally  
acceptable" includes residential and industrial uses.

25 Description of the Prior Art

Glyphosate (N-phosphonomethylglycine) well-known  
as an effective herbicide is an organic acid, and is  
relatively insoluble in water. Therefore, glyphosate is  
normally formulated and applied as a water-soluble salt,  
30 especially as the isopropylamine salt (IPA salt).

Various formulations of glyphosate are disclosed  
in U.S. Patents 4,405,531 issued to John E. Franz on  
September 20, 1983; 3,799,580 issued to John E. Franz on  
35 March 26, 1974; and 4,840,659 issued to John E. Franz on  
June 20, 1989. These patents are incorporated herein by  
reference in their entirety. Roundup® herbicide an  
aqueous concentrate comprising the IPA salt of

-2-

glyphosate is sold by Monsanto Company as an aqueous concentrate formulation which is normally diluted in water prior to application.

5           SharpShooter(TM) Herbicide Concentrate from Safer Inc. is said to contain fatty acid(s) as active ingredients such as those having 8 to 12 carbon atoms and mixtures thereof. It is believed used as a vegetation suppression agent.

10

SharpShooter(TM) is non-selective, and shows the result of its contact with weeds and unwanted vegetation in the form of necrosis, leaf burn, desiccation, wilting and the like. Typical fatty acids known to produce such  
15 symptoms are pelargonic and n-capric acid.

Pelargonic acid referred to also as nonanoic acid, nonylic acid and nonoic acid has the empirical formula  $C_9H_{18}O_2$ .

20

n-Capric Acid known also as n-decanoic acid, has the empirical formula  $C_{10}H_{20}O_2$ .

PCT/US88/03582 (W089/03178) of Safer, Inc.  
25 discloses a herbicidal composition and method for non-selectively controlling and retarding the growth rate and, if desired, causing extensive mortality of unwanted vegetation. The compositions consist essentially of one or more substances selected from the group consisting of  
30 aliphatic acids or their herbicidally active salts, disclosed preferably as octanoic acid, nonanoic acid, decanoic acid, n-decanoic acid, or dodecanoic acid, and an ammonium compound, preferably ammonium nitrate, sulfate or sulfamate. The composition is said to cause  
35 a plant mortality significantly in excess of the expected additive mortalities of the individual components.

-3-

U.S. Patent 4,975,110 issued to George S. Puritch et al discloses an environmentally compatible herbicidal composition, consisting essentially of a herbicidally effective amount of a saturated linear monocarboxylic fatty acid selected from the group consisting of the acids caprylic, pelargonic, capric, undecanoic, lauric and mixtures thereof; and a surfactant component.

U.S. Patent 3,870,503 issued to Louis G. Nickell on March 11, 1975 discloses that sucrose yield of sugarcane is increased by treating sugarcane a few weeks prior to harvest with a sugar cane ripening agent selected from the group consisting of n-valeric (pentanoic) acid and alkali metal salts or ethyl esters of an aliphatic monoacid having from one to five carbon atoms.

U.S. Patent 4,134,754 issued to Otto L. Hoffmann on January 16, 1979 discloses that the activity of barban is enhanced and variation of selectivity with climatic conditions is alleviated by applying to wild oats an effective amount of a composition comprising one part by weight carbon and at least four parts by weight of a polyunsaturated fatty acid exemplified by linolenic acid. The composition is preferably either dispersed in water with the aid of an emulsifier or dissolved in a mixture of water and a volatile organic solvent such as acetone.

U.S. Patent 4,436,547 issued to Michael J. Sampson on March 13, 1984 discloses that the effect of fungicides, herbicides, insecticides, nematocides and plant-growth regulators, is improved by co-administration of them with one or more of the following additives: carbohydrates, organic acids (particularly fatty acids and acids of the Krebs tricarboxylic acid cycle), vitamins and co-enzymes, purine and pyrimidine nucleosides and nucleotides, naturally occurring fats

-4-

and oils, certain amino acids and (but not where the agricultural chemical is itself a plant-growth regulator) plant-growth regulators. The invention is said to provide compositions containing one or more of the said agricultural chemicals and one or more of the said additives and methods of improving the harvest of a given crop by applying to it one or more of the said agricultural chemicals and one or more of the additives, either simultaneously or within up to about ten days of one another.

U.S. Patent 4,626,274 issued to Heinz Hausmann on December 2, 1986 discloses that a known herbicide such as a urea, carboxylic acid ester, aminoacid, benzoic derivative, benzonitrile, phenol derivative, diphenyl ether, triazinone, triazinedione, heterocycle, dipyridil derivative or benzosulphonamide, is rendered more effective by being combined with a synthetic spreading agent such as a silicone oil, etc.

20

US Patent 4,966,728 issued to James L. Hazen, on October 30, 1990 discloses herbicide adjuvants which are said to enhance the effectiveness of a broad spectrum of postemergent herbicides. These adjuvants preferably contain a low foaming nonionic surfactant, an anionic surfactant, a lower alkanol ester of a fatty acid, and a hydrocarbon oil component.

US Patent 4,902,334 issued to Azuma et al on February 20, 1990 discloses a plant metabolism regulating agent comprising as an active ingredient an alpha, beta or beta, gamma-unsaturated carboxylic acid or its derivative. This concept is said to be useful for controlling the metabolism of a plant, facilitating the growth of a beneficial plant such as cereals by inhibiting the growth of undesirable plants or eradicating them, regulating the growth of a plant and dwarfing a plant.

-5-

US Patent 4,904,645 issued to George S. Puritch et al on February 27, 1990 discloses a combination of pyrethrum and fatty acid salt material which is said to provide a stable, commercially useful and environmentally safe pesticidal formulation. The specific composition comprises an aqueous solution having a pH within the range of 7.5 to 8.8; an aqueous solution comprising about 50% by weight of monocarboxylic acids and their alkali metal salts, where the acid mixture is at least 70% oleic acid and 6% linoleic acid; a pyrethrum extract; a solvent for the pyrethrum, which is preferably a 2-6 carbon alcohol; a trace amount of an antioxidant. The composition is said to be effective against insects of at least the orders Homoptera, Coleoptera, Dermaptera, Hemiptera, and Lepidoptera, and against crustacea of the order Isopoda.

#### OBJECTS OF THE INVENTION

An objective of this invention is to provide an aqueous concentrate or ready to apply formulation comprising glyphosate or a water soluble salt of glyphosate and a fatty acid or a salt thereof or a mixture of fatty acids and salt thereof or a mixture of salts of the fatty acid(s) such that the early burndown symptoms on plants of the fatty acid and/or its salt or mixtures thereof are seen and the long term herbicidal control achieved by the use of the glyphosate is maintained and not sacrificed, eliminated or reduced by the contact action of the fatty acid or its water soluble salt. Enhanced shelf-stability is provided when the application pH is in the preferred range.

Without being bound by theory, it is believed that the fatty acid is the responsible moiety for providing the early contact visible symptoms on plants treated with a fatty acid salt contained in a formulation of this invention.

BRIEF DESCRIPTION OF THE INVENTION

The foregoing objects are provided in an agriculturally acceptable herbicidal composition  
5 comprising herbicidally effective amounts of each of:

- a) N-phosphonomethylglycine (glyphosate) or an agriculturally acceptable water soluble salt of N-phosphonomethylglycine or a compound which metabolizes  
10 thereto and
- b) a fatty acid such as a C<sub>5</sub> to C<sub>16</sub> or preferably a C<sub>8</sub> to C<sub>12</sub> saturated or unsaturated, straight or branched chain fatty acid or an agriculturally acceptable salt(s) thereof or a mixture of fatty acids  
15 and salt(s) thereof or a mixture of salts of a fatty acid(s) recognizing that such fatty acids may themselves be mixtures.

A preferred composition is wherein the ratio of  
20 a) to b) is in the range from about 1:10 to about 10:1 by weight and more preferably wherein the ratio of a) to b) is in the range from about 1:5 to about 5:1 and most preferably in the range from about 1:3 to about 3:1.

25 Especially preferred are the mono-isopropylamine and the trimethylsulfonium salts of N-phosphonomethylglycine.

Especially preferred are the potassium salts of  
30 capric fatty acid and pelargonic fatty acid and mixtures thereof. Potassium nonanoate and potassium decanoate may be employed.

Also disclosed is a method of killing or  
35 controlling plants where a herbicidally effective amount of the above described herbicidal composition is applied to said plants to kill or control said plants and wherein the early burndown effect of application of the



-7-

fatty acid or a salt thereof can be readily seen leaving intact the longer term control opportunity which is provided by simultaneous application of glyphosate or of the glyphosate moiety in the form of a water soluble  
5 salt of glyphosate or a compound which metabolizes to glyphosate.

Also disclosed is a process for preparing herbicidal compositions of this invention wherein a base  
10 such as those providing hydroxide in water preferably an amine or alkali metal or alkaline earth metal hydroxide most preferably potassium hydroxide is employed to control the desired final adjusted pH in the range from about 6.4 to about 7.8 and preferably in the range from  
15 about 6.8 to about 7.0 to provide an improved storage stable formulation, which is ready to use or ready to dilute.

#### Detailed Description of the Invention

20 The present invention relates to an agriculturally acceptable herbicidal, shelf-stable ready to use or shelf-stable aqueous concentrate formulation of N-phosphonomethylglycine or a water soluble salt of N-phosphonomethylglycine or salts or mixtures thereof,  
25 with one or more C<sub>5</sub> to C<sub>16</sub> or preferably C<sub>8</sub> to C<sub>12</sub> saturated or unsaturated, straight or branched chained fatty acid itself or in the form of a salt, salts or various mixtures thereof.

30 Various optional ingredients may also be employed if desired although not required. Greater or lesser amounts of those optional additaments may be employed as desired.

35 When the application pH of the formulation of the present invention is in the range from about 6.4 to about 7.8, and more preferably in the range from about 6.8 to about 7.0, the formulations of this invention

-8-

show increased shelf-stability in that they do not experience separation such as salting out or settling out of one or more ingredients when stored for prolonged periods of time at normal room temperatures or when  
5 undergoing accelerated aging studies done at 50°C or 60°C or at freezing temperatures.

As used herein, the phrase "early burndown symptoms" means visible plant symptoms characteristic of  
10 the contact action of a fatty acid or a salt thereof upon the leaf of a plant when applied in an effective amount alone to the surface of a plant. Such symptoms illustratively include necrosis, leafburn, desiccation, wilting and the like as visible evidence of herbicidal  
15 effectiveness.

As used herein, the term "long term control" means a significant inhibition of regrowth which is achieved through the use of a herbicidally effective  
20 amount of glyphosate (or glyphosate salt or a mixture thereof) on a plant to be killed or controlled.

The formulation of the present invention contains at least one water-soluble salt of glyphosate or a  
25 mixture of such salts. Several processes for the preparation of glyphosate and its water soluble salts are disclosed in the patent and chemical literature, e.g., U.S. Patents 3,977,860 and 4,486,358. Illustrative suitable water-soluble salts of glyphosate  
30 useful in the present invention are disclosed in U.S. Patent 4,405,531. The isopropylamine salt of glyphosate is most preferred in the formulation, preparation of the formulation and use of the present invention.

35 If desired, one or more of ammonium sulfate or potassium sulfate or any other compatible active or compatible nonactive ingredient may be employed as an additional ingredient(s) in a process for preparing a

-9-

formulation of the present invention, in its preparation and its use.

5 If ammonium sulfate is used during the formulation process, the formulation is normally filtered (or optional equivalent means) to remove any insoluble particulate materials which may be present in some commercial grades of ammonium sulfate.

10 The concentrates and solutions of the invention may contain optional additional components, for example anti-freeze agents such as ethylene glycol and propylene glycol.

15 Other examples are dyes, thickening agents, anti-foam agents, for instance silicone-based anti-foam agents, and surfactants, for instance non-ionic or cationic surfactants.

20 Additional illustrative suitable nonionic surfactants are recited in U.S. Patent 4,405 531 supra. Other suitable nonionic surfactants will be known to those skilled in the art.

25 Other surfactants which may be employed include alkylamine oxides, alkyl glucosides, ethoxylated or propoxylated quaternary amines and dialkylacetylene surfactants.

30 Particularly preferred, although optional, is an ethoxylated tallow amine surfactant having a degree of ethoxylation in the range from about 15 to about 18, mixtures thereof and the like. One such surfactant is Entry II, sold by Monsanto Company, St. Louis, Missouri  
35 USA.

The amount of surfactant optionally employed is typically in the range from about 0 to about 2 percent

-10-

by weight of a ready to use formulation of a spray solution formed by dilution with water of a concentrate formulation of this invention.

5           The concentrates and ready to use solutions of this invention may also be mixed with other water-soluble herbicides, for example but not limited to, salts of 2,4- dichlorophenoxyacetic acid, dicamba or 4-chloro-2-methylphenoxyacetic acid, or with finely-  
10 divided water-insoluble herbicides, for example but not limited to triazines, substituted ureas, sulfonylureas diphenyl ethers, dinitroanilines, pyridines and the like.

15           While Roundup® herbicide is the preferred source of a glyphosate water soluble salt, other suitable formulations providing a water soluble glyphosate salt or glyphosate acid may be employed if desired.

20           Alternatively, the water soluble glyphosate salt or an aqueous solution thereof may be prepared and then added directly as one of the process preparation steps.

          Some compositions of the invention were prepared  
25 by stirring about a 62 percent by weight aqueous solution of the mono-isopropylamine salt of glyphosate at about room temperature with the desired amount of fatty acid as a potassium salt and optionally, with a saturated solution of ammonium sulfate.

30           The formulations of the present invention may be conveniently prepared by mixing the desired ingredients together as in a blender or in any suitable container or device producing the needed amount of agitation  
35 resulting in mixed ingredients.

          Formulation examples were generally prepared by the following process:

-11-

- (1) adding all or a portion of the desired amount of initial water in a first container,
- (2) adding a portion of water and potassium hydroxide to the desired amount of fatty acid in a  
5 separate container to form a premix,
- (3) adding the fatty acid containing the potassium hydroxide in the premix from the separate container to the first container containing the initial water,
- 10 (4) adding a water soluble salt of N-phosphono-methylglycine alone with surfactant to the first container to form a desired final non pH adjusted mixture/solution,
- (5) adjusting the mixture/solution to a final  
15 desired pH by adding appropriate amounts of a concentrated potassium hydroxide solution and thereafter if desired,
- (6) admixing any remaining water and optional ingredients such as an anti-microbial for example  
20 Proxel® GXL (CAS 2634-33-5), sorbic acid, Legend MK, mixtures thereof and the like.

The conversion of a fatty acid such as pelargonic acid to salt such as a potassium salt is the step  
25 requiring most scrutiny of the process of this invention.

Pelargonic acid is a thin, clear oil at room temperature which sits on the surface of water when  
30 added. With mild agitation, the conversion time to form the salt of the fatty acid is slow. Extreme agitation and long batch stirring time are generally necessary to make the conversion complete which is desired.

35 In laboratory test batches, preparation of formulations of this invention was generally done by making the fatty acid salt in a separate smaller premix vessel using a beaker and vigorous agitation. This

-12-

premix was then added to the already mixed ingredients in a larger first container.

As used herein the term "vigorous agitation" includes stirring or mixing by any mechanical or equivalent means to produce a well mixed composition providing thorough blending.

The process of preparing a formulation of this invention may be done at temperatures in the range from about 10°C to about 100°C and preferably from about 20°C to about 40°C.

As another optional ingredient, an anti-foam agent may be included. Various suitable anti-foam agents include SAG 47 which is preferred as an anti-foam agent and Silicone Anti Foam® 30 IND.

The amount of anti-foam agent optionally employed is that which is sufficient to control and reduce an amount of foam which may be formed during the process of preparation of the formulation of this invention to a desired satisfactory level from a user's viewpoint. Generally the concentration of anti-foam agent is in the range from about 0.001% by weight of the total ingredients to about 0.1% although greater or lesser amounts of optional anti-foam agent may be employed.

Alternatively, a premix of about 1 part by weight Roundup® herbicide, about 2 parts water and potassium hydroxide as a 20 percent by weight solution was used to form a premix. Pelargonic acid was blended in this premix but thereafter separated easily. The benefit of this (alternative) process of preparation was that the premix went into the batch water with only mild agitation. Care must be given to prevent possible heavy residue in the premix beaker.

-13-

The most preferred previously described process requires some agitation. It may be likely that the addition of the fatty acid at the suction side of a flow through homogenizer Tekmar for example, into the  
5 potassium hydroxide composition would be sufficient to produce the desired amount of agitation.

In a preferred process for preparing formulations of this invention, the vessel wherein the formulation  
10 will be 50 gallons or more is initially charged with about 25% of the total desired water. Gentle agitation or recirculation is begun during and after this initial water addition. A base providing hydroxide in water is added as is optionally an ethoxylated tallowamine  
15 surfactant such as Entry II® Surfactant, sold by Monsanto Company, St. Louis, Missouri USA.

Agitation is increased and the added fatty acid is allowed to react. When the reaction is considered  
20 complete, the fatty acid appears homogeneous and not aggregated at the surface of the resulting composition. About half of the remaining desired water is added as are desired amounts of the isopropylamine salt of glyphosate, and optionally Proxel and Sag 47 with time  
25 allowed for adequate mixing. Thereafter remaining desired water is added and the pH is adjusted using potassium hydroxide.

Ready to use formulations having C<sub>9</sub> or C<sub>10</sub> fatty  
30 acids in the range from about 0.5 to about 1% by weight of the total spray weight were the most effective formulations from a herbicidal efficacy viewpoint with mixtures thereof providing similar efficacy.

35 The formulations of the present invention can be readily used as prepared or further diluted in water by a user in a spray container prior to use, although those

-14-

of skill in the art will recognize that the dilution will have a practical dilution limit set by efficacy.

Suitable herbicidally efficacious application rates of glyphosate will vary depending on plant species, weather, climate and geography. The spray volume is in the range from about 10 to about 1000 gallons per acre (gpa) (93.6 liters per hectare to about 9360 liters per hectare) preferably from about 50 to about 250 gpa (468 - 2340 liters per hectare). US Patent 3,799,758 provides illustrative application rates of glyphosate on an acid equivalent basis.

The following examples are presented to illustrate the present invention as well as some of the various embodiments of the invention. These examples are presented as being illustrative of the novel formulations and are not intended to be a limitation of the scope thereof. Rather these examples are intended to teach by illustration and provide what is believed to be a useful and readily understood description of procedures to make and use the invention.

In the examples, all parts, percentages and properties are by weight unless otherwise indicated. In some examples, comparative formulations illustrative of the prior art were made and compared to compositions of this invention.

### EXAMPLES

#### General Methods

Tall fescue, common bermuda, common dandelion, sweet clover and large crabgrass were grown from seed in 4 inch (10.16 centimeters) square pots containing a mixture of soil and commercial soil conditioners. After planting, the soil was kept moist and seedlings emerged in 5-8 days. Approximately 2 weeks after emergence, plants were mowed to a uniform height of 6 cm. Height



-15-

was maintained at 6 cm with 3 mowings per week for fescue, bermuda and crabgrass and with 1 mowing per week for dandelion and clover. Unless otherwise noted, all species except crabgrass were about 8 weeks old at the time of treatment. Crabgrass was 4-5 weeks old. Treatments were applied with a Devilbiss atomizer using compressed air. Chemicals were formulated as described for individual tests. For quick symptom development (1-3 DAT), evaluations were made on a 0-3 scale where

10

0=no effect

1=injury noticeable compared to untreated control

2=injury obvious

3=injury dramatic

15

Percent control ratings at later time periods were made on a 0-100 scale where 0=no effect and 100 is complete death. Results are the average of 2 replications.

20

#### Example 1

Combinations of C<sub>8</sub>-C<sub>12</sub> saturated fatty acids as its potassium salt with Glyphosate and an ethoxylated tallow amine surfactant having a degree of ethoxylation of about 15 to 18 were prepared.

Eight formulations containing either 0.5% or 1% C<sub>8</sub>, C<sub>9</sub>, C<sub>10</sub> or C<sub>12</sub> saturated fatty acids in 0.96% glyphosate-IPA and 0.36% ethoxylated tallow amine surfactant were applied respectively at 225, 112 or 56 gallons per acre (2106, 946, or 524 liters per hectare) to 8 week old fescue and bermuda, 5 week old dandelion and sweet clover and 4 week old crabgrass. Visual observations of burndown were made 1 DAT (Day After Treatment) on 0-3 scale and at 26 DAT on 0-100 scale. Results are in Table 1.

Table 1. Weed control with combinations of fatty acids, glyphosate and ethoxylated tallow amine surfactant.

Fatty acid	L/L	Fescue		Bermuda		Dandelion		Clover		Crabgrass	
		1*	26*	1	26	1	26	1	26	1	26
None } Prior	2106	0**	98***	0	98	0	100	0	70	0	98
None } Art	946	0	95	0	95	0	95	0	40	0	98
None }	524	0	80	0	70	0	98	0	30	0	95
1% C <sub>12</sub>	2106	0	98	0	98	0	98	2	60	2	99
1% C <sub>12</sub>	946	0	98	0	95	0	100	2	30	2	98
1% C <sub>12</sub>	524	0	85	0	70	0	95	0	30	1	98
0.5% C <sub>12</sub>	2106	0	99	0	98	0	100	2	80	2	100
0.5% C <sub>12</sub>	946	0	95	0	90	0	98	1	30	2	99
0.5% C <sub>12</sub>	524	0	80	0	60	0	95	0	20	1	95
1% C <sub>10</sub>	2106	2	98	1	95	2	100	3	80	3	98
1% C <sub>10</sub>	946	1	98	0	85	0	100	2	50	2	99
1% C <sub>10</sub>	524	0	90	0	50	0	75	2	30	1	95
0.5% C <sub>10</sub>	2106	2	100	1	95	2	100	2	60	3	98
0.5% C <sub>10</sub>	946	1	95	0	90	0	100	2	30	2	98
0.5% C <sub>10</sub>	524	0	90	0	70	0	95	1	30	1	95
1% C <sub>9</sub>	2106	2	98	2	95	2	100	3	70	3	98
1% C <sub>9</sub>	946	2	95	1	90	1	98	2	50	3	99
1% C <sub>9</sub>	524	1	80	0	60	0	95	2	30	1	95
0.5% C <sub>9</sub>	2106	2	99	1	95	1	100	2	60	2	98
0.5% C <sub>9</sub>	946	1	95	0	90	0	100	1	50	1	95
0.5% C <sub>9</sub>	524	0	80	0	60	0	85	0	20	1	95
1% C <sub>8</sub>	2106	2	99	2	95	1	100	2	60	3	98
1% C <sub>8</sub>	946	1	95	0	90	0	95	1	30	2	99
1% C <sub>8</sub>	524	0	80	0	50	0	90	1	20	0	98
0.5% C <sub>8</sub>	2106	1	99	1	95	0	100	2	70	2	99
0.5% C <sub>8</sub>	946	1	95	0	90	0	100	2	50	1	98
0.5% C <sub>8</sub>	524	0	70	0	50	0	95	0	30	0	98

\* Days after treatment

\*\* Rated on 0-3 scale

\*\*\* Rated on 0-100 scale

-17-

Example 2

Efficacy of C<sub>9</sub> and C<sub>10</sub> saturated fatty acids as the potassium salt alone and in mixtures in combinations with glyphosate.

Eight week old tall fescue and bermuda, 4 week old crabgrass and 2 week old yellow nutsedge were sprayed with C<sub>9</sub> and C<sub>10</sub> saturated fatty acids at 0.5 and 1% w/v alone and in mixtures at volumes of 56, 112 and 225 gal/A (524, 946 and 2106 liters per hectare). Yellow nutsedge had not been mowed prior to spraying. Visual observations of burndown were made on a 0-3 scale 1 DAT and percent control was evaluated on a 0-100 scale 29 DAT. Results are in Table 2.

In Table 2 following C<sub>9</sub> and C<sub>10</sub> fatty acids and mixtures of both give significant enhancement of final control of yellow nutsedge by glyphosate, as indicated. Mixtures appear more potent in this regard than either C<sub>9</sub> or C<sub>10</sub> fatty acid on its own. The fatty acids without glyphosate have zero effect on yellow nutsedge. Yellow nutsedge is a major problem weed and one that is typically difficult to control with glyphosate-based products.



Example 3

Effect of C<sub>9</sub> and C<sub>10</sub> saturated fatty acids as its potassium salt on glyphosate.

5

Formulations containing 0.96% IPA-glyphosate (.74% ae) and 0.4% ethoxylated tallow amine surfactant having a degree of ethoxylation of 15 to 18 were prepared as shown in Table 3a. In addition a concentrated formulation containing 16.5% IPA-glyphosate, 6.15% ethoxylated tallow amine surfactant having a degree of ethoxylation of 15 to 18 and 4.1% C<sub>9</sub> + 4.1% C<sub>10</sub> fatty acid was diluted and sprayed at the same final concentration of active. Treatments were applied at 56 and 112 gal/A (524 and 946 liters per hectare). Visual observations of injury were recorded 1, 3 and 25 days after treatment. Results are shown in Tables 3a and 3b.

20

Example 4

Use of C<sub>9</sub> saturated fatty acid for enhanced burndown with Roundup® LLG ready-to-use herbicide.

Results of treatments of Roundup® LLG ready-to-use herbicide alone and in mixture with C<sub>9</sub> saturated fatty acid at 0.5% and at 1.0% of the spray solution are shown in Table 4.

30

Example 5

The effect of pH adjustment on performance of a mixture of Roundup® LLG ready-to-use herbicide containing 1% C<sub>9</sub> saturated fatty acid is shown in Table 5.

Table 3a. Glyphosate combined with saturated fatty acids applied at 56 gal/acre (524 liters per hectare).

	$C_p$	$C_{10}$	(Prior art)	Fescue			Clover			Dandelion		
				1*	3*	25*	1	3	25	1	3	25
0	0	0	0	0	0	85	0	1	50	0	0	70
.5%		0		0	1	80	2	2	50	0	0	70
1%		0		0	1	60	2	2	40	0	0	30
0		.5%		0	1	70	2	2	30	0	0	80
.5%		.5%		0	1	60	2	2	30	0	0	30
.5%**		.5%**		0	2	80	2	2	30	0	0	50

	$C_p$	$C_{10}$	(Prior art)	Bermuda			Crabgrass		
				1*	3*	25*	1	3	25
0	0	0	0	0	0	60	0	1	85
.5%		0		0	0	60	0	1	90
1%		0		0	0	50	0	1	90
0		.5%		0	0	50	1	2	80
.5%		.5%		0	1	50	1	2	70
.5%**		.5%**		0	0	40	0	1	70

\* Days after treatment  
 \*\* Diluted from concentrate

Table 3b. Glyphosate combined with saturated fatty acids applied at 112 gal/acre (946 liters per hectare).

C <sub>7</sub>	C <sub>10</sub>	(Prior art)	Fescue			Clover			Dandelion		
			1*	3*	25*	1	3	25	1	3	25
0	0	0	0	0	95	1	2	20	0	0	90
.5%	0	0	1	1	98	2	2	20	0	0	80
1%	0	0	1	1	98	2	2	20	0	0	80
0	.5%	.5%	1	2	98	2	2	30	0	1	85
.5%	.5%	.5%	1	2	99	1	2	40	0	1	60
.5%**	.5%**	.5%**	1	2	98	2	2	30	0	0	40

C <sub>7</sub>	C <sub>10</sub>	(Prior art)	Bermuda			Crabgrass		
			1*	3*	25*	1	3	25
0	0	0	0	0	80	1	2	95
.5%	0	0	0	0	80	1	2	95
1%	0	0	0	0	80	1	2	98
0	.5%	.5%	0	1	85	1	2	98
.5%	.5%	.5%	0	1	70	1	2	98
.5%**	.5%**	.5%**	0	0	80	1	2	90

\* Days after treatment  
 \*\* Diluted from concentrate

Table 4. Effect of C<sub>9</sub> fatty acid on performance of commercially available Roundup RTU herbicide.

Additive	Prior art	L/Ha	Fescue			Bermuda			Dandelion			Clover			Crabgrass		
			1*	20*	80***	1	20		1	20		1	20		1	20	
None	None	524	0**	80***		0	85		0	70		0	60		0	90	
None	None	946	0	95		0	98		0	100		1	90		0	98	
None	None	2106	0	99		0	98		0	100		1	95		0	100	
0.5% C <sub>9</sub>		524	0	90		0	80		0	80		2	50		0	95	
0.5% C <sub>9</sub>		946	0	98		0	90		0	100		2	85		1	95	
0.5% C <sub>9</sub>		2106	1	99		0	98		0	100		2	85		1	100	
1% C <sub>9</sub>		524	0	60		0	60		0	99		2	60		1	90	
1% C <sub>9</sub>		946	0	90		0	95		1	98		2	85		2	95	
1% C <sub>9</sub>		2106	1	95		0	95		1	98		2	95		3	99	

\* Days after treatment

\*\* Rated on a 0-3 scale

\*\*\* Rated on a 0-100 scale



Table 5. Effect of pH on activity of glyphosate with 1% C<sub>9</sub> fatty acid.

Treatment	L/Ha	Fescue			Clover			Dandelion		
		1*	2*	28*	1	2	28	1	2	28
Roundup RTU	524	0	0	0	0	1	0	0	0	60
	946	0	0	20	0	1	0	0	0	100
	1572	0	0	20	1	1	0	0	0	99
+FA** pH 6.4	2106	0	0	50	1	2	60	0	0	98
	524	0	1	10	2	2	0	0	0	50
	946	0	2	60	2	3	90	0	1	70
+FA** pH 6.75	1572	1	2	70	3	3	95	0	1	40
	2106	1	3	90	3	3	10	1	2	100
	524	0	1	10	2	2	30	0	0	50
+FA** pH 7.25	946	0	2	10	2	3	70	0	0	100
	1572	0	2	30	2	3	90	0	1	100
	2106	0	3	50	3	3	30	0	1	30
+FA** pH 7.71	524	0	0	0	2	2	0	0	0	50
	946	0	0	0	2	2	10	0	0	50
	1572	0	1	20	2	3	80	0	1	100
+FA** pH 7.71	2106	0	2	10	2	3	90	0	1	50
	524	0	0	0	2	2	50	0	1	50
	946	0	1	10	2	3	80	0	1	100
	1572	0	1	20	2	3	90	0	1	100
	2106	0	1	50	2	3	70	0	1	100

\* Days after treat

\*\* Roundup TRU + 1% C<sub>9</sub> fatty acid

Treatment	L/Ha	Bermuda			Crabgrass		
		1*	3*	28*	1	3	28
Roundup RTU	524	0	0	10	0	0	95
	946	0	0	70	0	0	95
	1572	0	0	70	0	0	95
	2106	0	1	90	1	1	95
+FA** pH 6.4	524	0	0	10	0	1	90
	946	0	1	60	1	2	98
	1572	0	2	60	1	3	98
	2106	0	2	90	1	3	98
+FA** pH 6.75	524	0	0	10	0	0	80
	946	0	1	50	1	2	95
	1572	0	1	50	1	2	95
	2106	0	1	50	1	3	95
+FA** pH 7.25	524	0	0	20	0	0	80
	946	0	1	20	1	2	99
	1572	0	1	90	1	2	98
	2106	0	2	90	1	3	98
+FA** pH 7.71	524	0	0	10	0	0	80
	946	0	1	10	0	1	95
	1572	0	1	10	0	2	95
	2106	0	1	50	0	2	95

\* Days after treatment

\*\* Roundup RTU + 1% C<sub>9</sub> fatty acid

-25-

Although this invention has been described with respect to specific embodiments, the details hereof are not to be construed as limitations, for it will be apparent that various equivalents, changes and  
5 modifications may be resorted to without departing from the spirit and scope of this invention and it is understood that such equivalent embodiments are intended to be included within the scope of this invention.

-26-

## WHAT IS CLAIMED IS:

1. An agriculturally acceptable herbicidally effective composition providing initial visible symptoms  
5 of contact of a fatty acid or fatty acid salt component hereof and longer term herbicidal kill or control by N-phosphonomethylglycine or an N-phosphonomethylglycine salt component hereof, comprising effective amounts of each of:
  - 10 a) a herbicidally effective amount of N-phosphonomethylglycine or one or more agriculturally acceptable water soluble salts thereof or mixtures thereof or a compound which metabolizes thereto and
  - b) a C<sub>5</sub> to C<sub>16</sub> saturated or unsaturated,  
15 straight or branched chain fatty acid or agriculturally acceptable salt thereof or mixture of fatty acids and salt(s) thereof or mixture of salts of fatty acid(s) with or without dilution in water having an application pH in the range from about 6.4 to about 7.8.  
20
2. The composition of Claim 1, wherein said fatty acid is in the range from C<sub>8</sub> to C<sub>12</sub> saturated or unsaturated, straight or branched chain fatty acid or agriculturally acceptable salt thereof or mixture of  
25 fatty acids and salt(s) thereof or mixture of salts of fatty acid(s) with or without dilution in water having an application pH in the range from about 6.4 to about 7.8.
3. The composition of Claim 1 wherein the ratio  
30 of component a) to component b) is in the range from about 1:10 to about 10:1.
4. The composition of Claim 2 wherein the ratio  
35 of component a) to component b) is in the range from about 1:5 to about 5:1.

-27-

5. The composition of Claim 2 wherein said agriculturally acceptable water soluble salt of N-phosphonomethylglycine is the isopropylamine salt of N-phosphonomethylglycine.

5

6. The composition of Claim 2 wherein said agriculturally acceptable water soluble salt of N-phosphonomethylglycine is the trimethylsulfonium salt of N-phosphonomethylglycine.

10

7. The composition of Claim 2 wherein said fatty acid salt is the potassium salt of capric acid or the potassium salt of pelargonic acid or a mixture of the potassium salts of capric and pelargonic acids.

15

8. The composition of Claim 4 wherein the pH of the composition after dilution in water if necessary is in the range from about 6.8 to about 7.0 and enhanced shelf-stability is provided.

20

9. The composition of Claim 6, wherein an additional surfactant is employed.

10. The composition of Claim 9, wherein said surfactant is an ethoxylated tallow amine.

25

11. The composition of Claim 9, wherein an additional antifoam is employed.

12. The composition of Claim 11 wherein the said antifoam is SAG 47.

30

13. A method of killing or controlling weeds where a herbicidally effective amount of the composition of Claim 1 is applied to said weeds to kill or control.

35

14. The method of Claim 13 wherein said composition is the composition of Claim 2.

-28-

15. The method of Claim 13 wherein said composition is the composition of Claim 3.

5 16. The method of Claim 13 wherein said composition is the composition of Claim 4.

17. The method of Claim 13 wherein said composition is the composition of Claim 5.

10

18. The method of Claim 13 where in said composition is the composition of Claim 6.

19. A process for preparing said composition of  
15 Claim 1 by adding together in any order

a) a herbicidally effective amount of N-phosphonomethylglycine or an agriculturally acceptable water soluble salt thereof or mixtures thereof and

b) a C<sub>5</sub> to C<sub>16</sub> or preferably C<sub>8</sub> to C<sub>12</sub> saturated  
20 or unsaturated, straight or branched chain fatty acid or an agriculturally acceptable salt(s) thereof or mixtures thereof, and wherein a base is employed to provide a said composition with a pH after dilution in water if necessary in the range from about 6.4 to about 7.8.

25

20. The process of Claim 19 wherein said pH is in the range from about 6.8 to about 7.0.

21. The process of Claim 20 wherein said pH is  
30 controlled by addition of a hydroxide or an amine.

22. The process of Claim 21 wherein said hydroxide is an amine, an alkali metal or an alkaline earth metal hydroxide.

35

23. The process of Claim 22 wherein said alkali metal hydroxide is potassium or sodium hydroxide.

-29-

24. The process of Claim 23 wherein said alkali metal hydroxide is potassium hydroxide.

25. A process of preparing an enhanced pesticide composition of Claim 1 by the additional step in any order of adding all or a portion of the desired amount of optional initial water in a first container.

26. The process of Claim 25 for preparing the enhanced pesticide composition of Claim 1 by the additional step of adding potassium hydroxide to the desired amount of fatty acid in a separate container from a first container to form a premix.

27. The process of Claim 26 for preparing the enhanced pesticide composition of Claim 1 comprising the further additional step of adding said fatty acid containing said hydroxide in the premix from the separate container to the one container containing the initial water and additional surfactant.

28. The process of Claim 27 further comprising the additional process step of adding the glyphosate salt, antimicrobial and antifoam to the mixture claimed in claim 27.

29. The process of Claim 28 which comprises the yet additional process step of adjusting said mixture/solution in claim 28 to a final desired pH by adding appropriate amounts of a hydroxide solution.

30. The process of Claim 29 which comprises the yet further additional process step of admixing any remaining water and optional ingredients to said composition of Claim 29.

# INTERNATIONAL SEARCH REPORT

International Application No PCT/US 91/09621

<b>I. CLASSIFICATION OF SUBJECT MATTER</b> (If several classification symbols apply, indicate all) <sup>6</sup> According to International Patent Classification (IPC) or to both National Classification and IPC IPC5: A 01 N 57/20, 37/00/(A 01 N 37/00, 37:02, 37:06)														
<b>II. FIELDS SEARCHED</b> <div style="text-align: center; border-top: 1px solid black; border-bottom: 1px solid black;">Minimum Documentation Searched<sup>7</sup></div> <table style="width: 100%; border-collapse: collapse;"> <tr> <th style="width: 25%; border-bottom: 1px solid black;">Classification System</th> <th style="border-bottom: 1px solid black;">Classification Symbols</th> </tr> <tr> <td style="padding: 5px;">IPC5</td> <td style="padding: 5px;">A 01 N</td> </tr> </table> <div style="text-align: center; border-top: 1px solid black; border-bottom: 1px solid black;">Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in Fields Searched<sup>8</sup></div>			Classification System	Classification Symbols	IPC5	A 01 N								
Classification System	Classification Symbols													
IPC5	A 01 N													
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT<sup>9</sup></b> <table style="width: 100%; border-collapse: collapse;"> <tr> <th style="width: 10%; border-bottom: 1px solid black;">Category *</th> <th style="border-bottom: 1px solid black;">Citation of Document,<sup>11</sup> with indication, where appropriate, of the relevant passages<sup>12</sup></th> <th style="width: 10%; border-bottom: 1px solid black;">Relevant to Claim No.<sup>13</sup></th> </tr> <tr> <td style="text-align: center; vertical-align: top; padding: 5px;">X</td> <td style="padding: 5px;">US, A, 4436547 (M.J. SAMPSON) 13 March 1984, see the abstract; column 2, lines 19-25; column 11, lines 10-29; column 4, lines 62-63</td> <td style="text-align: center; vertical-align: top; padding: 5px;">1-30</td> </tr> <tr> <td style="text-align: center; vertical-align: top; padding: 5px;">X</td> <td style="padding: 5px;">STN International, File CABA, STN accession no. 90:134023, A.J. Wells et al: "adjuvants, glyphosate efficacy and post-spraying rainfall", Plant Protection Quarterly, (1989) 4(4) 158-164.</td> <td style="text-align: center; vertical-align: top; padding: 5px;">1-30</td> </tr> <tr> <td style="text-align: center; vertical-align: top; padding: 5px;">X</td> <td style="padding: 5px;">STN International, File CABA, STN accession no. 91:10805, D.G. Shilling et al: "Influence of surfactants and additives on phytotoxicity of glyphosate to torpedograss", Journal of Aquatic Plant Management, (1990) 28 23-27</td> <td style="text-align: center; vertical-align: top; padding: 5px;">1-30</td> </tr> </table>			Category *	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>	X	US, A, 4436547 (M.J. SAMPSON) 13 March 1984, see the abstract; column 2, lines 19-25; column 11, lines 10-29; column 4, lines 62-63	1-30	X	STN International, File CABA, STN accession no. 90:134023, A.J. Wells et al: "adjuvants, glyphosate efficacy and post-spraying rainfall", Plant Protection Quarterly, (1989) 4(4) 158-164.	1-30	X	STN International, File CABA, STN accession no. 91:10805, D.G. Shilling et al: "Influence of surfactants and additives on phytotoxicity of glyphosate to torpedograss", Journal of Aquatic Plant Management, (1990) 28 23-27	1-30
Category *	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>												
X	US, A, 4436547 (M.J. SAMPSON) 13 March 1984, see the abstract; column 2, lines 19-25; column 11, lines 10-29; column 4, lines 62-63	1-30												
X	STN International, File CABA, STN accession no. 90:134023, A.J. Wells et al: "adjuvants, glyphosate efficacy and post-spraying rainfall", Plant Protection Quarterly, (1989) 4(4) 158-164.	1-30												
X	STN International, File CABA, STN accession no. 91:10805, D.G. Shilling et al: "Influence of surfactants and additives on phytotoxicity of glyphosate to torpedograss", Journal of Aquatic Plant Management, (1990) 28 23-27	1-30												
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p><b>* Special categories of cited documents:<sup>10</sup></b></p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance, the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance, the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&amp;" document member of the same patent family</p> </div> </div>														
<b>IV. CERTIFICATION</b> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%; border-bottom: 1px solid black; padding: 5px;">           Date of the Actual Completion of the International Search  <b>8th April 1992</b> </td> <td style="width: 50%; border-bottom: 1px solid black; padding: 5px;">           Date of Mailing of this International Search Report  <b>13.05.92</b> </td> </tr> <tr> <td style="border-bottom: 1px solid black; padding: 5px;">           International Searching Authority  <b>EUROPEAN PATENT OFFICE</b> </td> <td style="border-bottom: 1px solid black; padding: 5px;">           Signature of Authorized Officer  <div style="display: flex; align-items: center;"> <div style="margin-right: 20px;">Nicole De Blé</div> </div> </td> </tr> </table>			Date of the Actual Completion of the International Search <b>8th April 1992</b>	Date of Mailing of this International Search Report <b>13.05.92</b>	International Searching Authority <b>EUROPEAN PATENT OFFICE</b>	Signature of Authorized Officer <div style="display: flex; align-items: center;"> <div style="margin-right: 20px;">Nicole De Blé</div> </div>								
Date of the Actual Completion of the International Search <b>8th April 1992</b>	Date of Mailing of this International Search Report <b>13.05.92</b>													
International Searching Authority <b>EUROPEAN PATENT OFFICE</b>	Signature of Authorized Officer <div style="display: flex; align-items: center;"> <div style="margin-right: 20px;">Nicole De Blé</div> </div>													



III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
A	WO, A1, 9007275 (MONSANTO COMPANY) 12 July 1990, see the claims —	1-30
A	US, A, 4975110 (G.S. PURITCH ET AL) 4 December 1990, see the claims —	1-30
A	WO, A1, 8903178 (SAFER, INC.) 20 April 1989, see the claims —	1-30
P,A	WO, A1, 9105472 (SAFER, INC.) 2 May 1991, see the claims —	1-30
A	EP, A1, 0192583 (RHONE-POULENC AGROCHIMIE) 27 August 1986, see the claims —	1-30
A	DE, A1, 3247050 (BAYER AG) 20 June 1984, see claim 1, (I)(3)(a), (II)(2) — —————	1-30

**ANNEX TO THE INTERNATIONAL SEARCH REPORT  
ON INTERNATIONAL PATENT APPLICATION NO.PCT/US 91/09621**

SA 55900

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.  
The members are as contained in the European Patent Office EDP file on 28/02/92  
The European Patent office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A1- 9007275	12/07/90	AU-D- 4833390	01/08/90
		CA-A- 2006816	30/06/90
		EP-A- 0378985	25/07/90
		EP-A- 0452366	23/10/91
US-A- 4975110	04/12/90	WO-A- 91/05471	02/05/91
WO-A1- 8903178	20/04/89	AU-D- 2625288	02/05/89
		EP-A- 0335961	11/10/89
WO-A1- 9105472	02/05/91	US-A- 5035741	30/07/91
EP-A1- 0192583	27/08/86	AU-B- 577658	29/09/88
		AU-D- 5259586	31/07/86
		CA-A- 1272890	21/08/90
		FR-A-B- 2576181	25/07/86
		GB-A-B- 2169806	23/07/86
		JP-A- 61172805	04/08/86
		QA-A- 8194	30/10/87
DE-A1- 3247050	20/06/84	AU-D- 2237983	28/06/84
		EP-A-B- 0113857	25/07/84
		JP-A- 59118701	09/07/84
		US-A- 4626274	02/12/86

For more details about this annex : see Official Journal of the European patent Office, No. 12/82

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets

(11) Veröffentlichungsnummer:

(11) Publication number:

(11) Numéro de publication:

**0 566 648**

Internationale Anmeldung veröffentlicht durch die  
Weltorganisation für geistiges Eigentum unter der Nummer:

**WO 92/11764** (art.158 des EPf).

International application published by the World  
Intellectual Property Organisation under number:

**WO 92/11764** (art.158 of the EPC).

Demande internationale publiée par l'Organisation  
Mondiale de la Propriété sous le numéro:

**WO 92/11764** (art.158 de la CBE).

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
Y	Chemical Abstracts, volume 73, no. 2, 13 July 1970, (Columbus, Ohio, US), Nal'kina, Z.A. et al. : "Extraction of ammonium in the water-ammonium nitrate-ammonium caprylate-caprylic acid system. ", see page 305, abstract 7872b, & Inv.Sib.Old.Akad.Nauk SSSR, Ser.Khim.Nauk 1970, 1, 51- 7 --	1-8
Y	DE, A1, 28 48 184 (RALSTON PURINA CO.) 10 May 1979, see claims 1, 4, 7, 10 -- -----	1-8

**ANNEX TO THE INTERNATIONAL SEARCH REPORT  
ON INTERNATIONAL PATENT APPLICATION NO. PCT/US 88/03582**

SA 25219

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.  
The members are as contained in the European Patent Office EDP file on 12/01/89  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB-A- 1 243 987	25/08/71	NONE	
DE-A- 1 792 119	13/01/72	NL-A- 6810500	28/01/69
		FR-A- 1581109	12/09/69
		GB-A- 1237278	30/06/71
		US-A- 3619168	09/11/71
US-A- 2 277 744	31/03/42	NONE	
DE-A1- 28 48 184	10/05/79	FR-A-B- 2408310	08/06/79
		US-A- 4225621	30/09/80
		CA-A- 1119878	16/03/82